arXiv:1611.07003v2 [astro-ph.SR] 3 Dec 2016

Catalog of averaged magnetic phase curves of stars

V.D. Bychkov[1], L.V. Bychkova[1], J. Madej[2]
[1] Special Astrophysical Observatory, Russian Academy of Sciences
[2] Warsaw University Observatory, Poland

Abstract

The second version of the catalog contains information about 275 stars of different types. During the time that has elapsed since the creation of the first catalog, situation fundamentally changed primarily due to the significant increase of accuracy of magnetic fields (MF) measurements. Up to now global magnetic field were discoverd and measured in stars of many types and their behavior partially was studied. Magnetic behavior of Ap and Bp stars is the most thoroughly studied. The catalog contains information about 182 such objects. The main goals for the construction of the catalog are: 1. Review and summarize our kowledge about magnetic behavior of different types of stars. 2. The whole data are uniformly presented and processed which will allow one to perform statistical analysis of the variability of the longitudinal magnetic fields of stars. 3. The informations are presented in the most convenient form for testing different theoretical models of different kind. 4. The catalog will be useful for the development of observational programs.

submitted to

Stars: from collapse to collapse
Special Astrophysical Observatory
2016 October 3-7, Nizhnij Arkhyz
Karachai-Cherkessian Republic (Russia)

1 Introduction

We present and analyze results of the longitudinal magnetic field B_e measurements of stars collected from different bibliographic sources and personal communications, as well as our own determinations obtained at Special Astrophysical Observatory (Russian Academy of Sciences), see Bychkov (2008) and Bychkov et al. (2012).

Figs. 1-4 present sample rotational magnetic phase curves (MPC) for stars of various types and the distribution of the available MPC over spectral classes.

2 Description of the catalog

For each star in the catalog the average magnetic phase curve was obtained by the least squares fitting of a sine wave or a double wave to observed B_e points. This is the same method as that used in the first version of the catalog (Bychkov et al. 2005). In general the second version of the catalog is mostly very similar to the first version.

Recently the amount of available data sharply increased partly due to the flow of new high-precision observational data obtained with the Least Squares Deconvolution method (LSD) described by Donati et al. (1997). These circumstances caused discovery of variable longitudinal magnetic fields in a number of stars of different types and more precise determination of MPC's in magnetic stars already known.

Fig. 1 shows the distribution of stars with known MPC among spectral types. As the figure shows, the bulk of stars for which MPC's are known is concentrated in the spectral region around type A. This is due to the large contribution from Ap/Bp stars.

Table 1 shows the types of stars for which MPC's were determined. Precise specification of types of stars sometimes is not possible. For example, HD96446 consists of a He-r and β Cep stars, HD97048 consists of TTS (T Tau) and Ae/Be Herbig objects, flaring DT Vir consists of UV Cet + RS CVn types (Flare + RS CVn type stars) .

The best investigated stars in the Ap/Bp group, in turn, can be divided into two subclasses. The first subclass consists of stars who have simple sine wave MPC (total 141 objects). Fig. 4 shows the distribution of rotational periods in this group of stars. Stars with long periods of rotation higher than the maximum period in Fig. 4 are: HD2453 period 520.5 d, HD5797 – 68.046 d, HD8441 – 69.2 d, HD37058 – period 223.75 d HD41403 – 4750. d, HD55719 – 740.247 d, HD187474 – 2345.8 d, HD188041 - 222.63 and HD201601 – 35462.5 days.

Fig. 5 shows the distribution of coefficiens B_0 (average field strength) and B_1 (half amplitude) of the best sine wave fits for stars with simple



Figure 1: Left: Distribution of stars with known magnetic phase curves over spectral types. Right: Magnetic phase curve of the Ofp? star HD191612 with a rotational period of 537.6 days (Wade et al. 2011), the example of supermassive supergiants of the Ofp? type.



Figure 2: Left:Classical T Tau star (CTTS) V2129 Oph Magnetic phase curve for the T-Tauri type star V2129 Oph. The period of rotation equals 6.53 days, according to Donati et al. (2007). Right:Magnetic phase curve for chemically normal stars HD149438, assuming the rotational period of 41.033 days from Donati et al. (2006).

MPC.

The second group of Ap/Bp stars form objects whom IFC are double sine wave. This group consists of 45 stars. Fig. 6 shows the distribution of rotational periods in this group. Stars with extremely long periods did not fit into Fig. 6: these are HD965 – period 7125.65 d, HD9996 – 7961.8 d, HD18078 – 1358. d, HD94660 – 2800.0 d and HD126515 – 129.95 days.

Figs. 6-7 depict distribution of the coefficients B_0, B_1 and B_2 of double sine waves best fitted to the magnetic B_e time series observed in these stars. In the first version of the catalog (Bychkov et al. 2005) we pointed to the

Table 1: Number of objects of various types with known MPC

Ap/Bp	182	TTS	3
var.beta Cep type	14	var.Ori type	2
Flare stars	12	host Planet star	2
multiple star	11	Be-star	2
Ae/Be	9	semi-reguliar var.pulsating	2
Rotationally var.star	7	Hosts a planetary system	2
SPBS type	6	var.delta Cep type	1
HPMS	5	Pulsating star	1
var.BY Dra	5	var.type delta Sct	1
Of	4	DA	1
normal chem.comp.stars	4	WN D	1



Figure 3: Left:Magnetic phase curve of the Ae/Be Herbig star HD101412 with the rotational period of 42.076 days by Hubrig et al. (2011), see also Jarvinen et al. (2016). Right:Magnetic rotational phase curve for the mCP star HD 32633 obtained from highly accurate observations by Silvester et al. (2012).

unusually large bias of B_0 towards negative values. We attributed the effect to the low statistics of total 19 double wave stars in the old catalog. Actually we have 45 stars in the sample, over twice more than before. Now the distribution of B_0 is more symmetric around the value of zero and only a slight imbalance towards the negative values of the average B_0 still remains there.

Only in one exceptional star HD37776, the MPC is best described by the triple sinusoid.



Figure 4: Left: Magnetic rotational phase curve for the SPBS star HD 3360 obtained from highly accurate observations by Briquet et al. (2016). Right:Distribution of periods for Ap stars with simple harmonic magnetic phase curves.



Figure 5: Left: Distribution of the parameter B_0 for Ap stars with simple MPC. Right:Distribution of the halfamplitude B_1 for Ap stars with simple MPC.

3 Conclusions

Catalog of the magnetic rotational phase curves MPC primarily represents a review and synthesis of the longitudinal magnetic field measurements of stars of various classes, accumulated up to now. Principal reason of the longitudinal field variability is rotation of a star with its magnetic field frozen into ionized gas and seen at different aspect angles.

Magnetic field itself evolves relatively rapidly, which can be observed in cool red dwarfs (flare stars). This is a consequence of the strongly differential rotation of those stars, which causes steady generation of local magnetic fields, which sometimes annihilate one another causing flares. On the con-



Figure 6: Left:Distribution of periods for Ap stars with complex MPC. Right:Distribution of the parameter B_0 for Ap stars with complex MPC.



Figure 7: Left:Distribution of the halfamplitude B_1 for Ap stars with complex MPC. Right:Distribution of the halfamplitude B_2 for Ap stars with complex MPC.

trary, sometimes local fields add up to change strength and configuration of the global magnetic field in time scales from months to years. This causes changes of parameters and shape of the magnetic phase curve.

In case of stars of other types, global magnetic fields do not evolve rapidly. Nevertheless, there are apparent indications that in some stars there are signatures of rapid evolution of the global magnetic fields.

We collected and uniformly processed all available observations on the variability of longitudinal magnetic fields of stars in the unified form, which allows one for a statistical analysis of the data and for testing and verification of various theoretical models. The catalog will be also useful for the development of observational programs. We believe, that the accumulated observational data will be used to refine rotational periods and parameters

of many stars included in this catalog.

4 Acknowledgements

This work was supported by the Russian Science Foundation (project No. 14-50-00043).

References

[1] Briquet, M., Neiner, C., Petit, P., Leroy, B., de Batz, B., 2016, A&A, 587, A126

[2] Bychkov, V.D., Bychkova, L.V., Madej, J. 2005, A&A, 430, 1143

[3] Bychkov, V.D., Bychkova, L.V., Madej, J. 2016, in press

[4] Bychkov, V.D., Bychkova, L.V., Madej, J., Panferov, A.A. 2015, ASP conf. ser., 494, 210

[5] Bychkov V.D., Bychkova L.V., Madej J., Topilskaya G.P. 2016, arXiv:1606.09562

[6] Donati, J.F., Semel, M., Carter, B.D., Rees, D.E., Cameron, A.C. 1997, MNRAS, 291, 658

[7] Donati, J.-F., Howarth, I.D., Jardine, M.M., et al. 2006, MNRAS, 370, 629

[8] Donati, J.-F., Jardine, M.M., Gregory, S.G., et al. 2007, MNRAS, 380, 1297

[9] Donati, J.-F., Morin, J., Petit, P. et al. 2008, MNRAS, 390, 545

[10] Hubrig, S., Mikulasek, Z., Gonzalez, J.F., et al. 2011, A&A, 525, L4

[11] Jarvinen, S.P., Hubrig, S., Scholler, M., Ilyin, I., Carroll, T.A., Korhonen, H., 2016, AN, 337, 329

[12] Silvester, J., Wade, G.A., Kochukhov, O., et al. 2012, MNRAS, 426, 1003

[13] Wade, G.A., Bagnulo, S., Drouin, D., Landstreet, J.D., Monin, D., 2007, MNRAS, 376, 1145

[14] Wade, G.A., Howarth, I.D., Townsend, R.H.D., et al. 2011, MNRAS, 416, 3160